



REPULS®

REPULS® Voluntary Exposure Checklist

Manufacturer's Recommended Procedures

AUTHORIZED USE

Voluntary exposure procedure is authorized using MK3/MK4 canister of REPULS. It is the REPULS Instructor's responsibility to ensure completion of all steps below.

PRE-EXPOSURE PREPARATION

- Ensure the REPULS Voluntary Exposure Waiver is complete and signed by the individual to be sprayed. Waiver: <https://forms.office.com/r/vSTeq21aUG>
- Confirm each individual is fit for duty with their organization prior to participating.
- Test spray the canister with a quick burst to confirm it is functioning properly.
- Confirm the training area is safe and free from hazards.
- Have a safety person standby with two clean towels, at minimum one 16–20 oz bottle of water, and sterile saline solution or eye wash.
- Brief the individual on the exposure and post-exposure treatment process.

EXPOSURE PROCEDURE

IMPORTANT

REPULS does not contain OC or CS, does not cause cross-contamination, and does not reactivate. REPULS and pepper spray of any brand shall NOT be deployed in close proximity. REPULS shall not be sprayed directly into someone's eyes at less than 3 feet. Consult a physician if the individual has had recent LASIK or eye surgery. Crotega does not assume liability for product misuse.

- Deploy REPULS from approximately 8–10 feet using a current MK3/MK4 REPULS canister.
- Instruct the individual to close their eyes and not open them until instructed. Confirm the individual is ready.
- Spray the individual's forehead only — soaking the forehead. Tell the individual to open their eyes, then immediately instruct them to keep their eyes closed to relieve discomfort.
- Do NOT have the individual try to forcibly open their eyes or perform tasks.
- Do NOT have the individual rub their eyes.

POST-EXPOSURE TREATMENT

- Begin post-exposure treatment immediately. Do not delay — delays may increase the risk of an adverse reaction.
- Pour water starting at the individual's forehead. Gently wipe the face with a clean towel, keeping eyes closed to avoid scratching.
- Alternate pouring water into each eye; ask the individual to blink while pouring. Use sterile saline solution at least twice on each eye.
- After using 2–3 oz of water, apply saline solution or eye wash to both eyes and gently wipe with a clean towel. Continue alternating saline solution or eye wash and water until the entire bottle of water is used.
- Remain with the individual for a reasonable period of time after symptoms subside to monitor for adverse reactions or eye irritation.

MEDICAL

Anyone sprayed with REPULS should receive medical attention if they request it or if symptoms or eye irritation persist after 30 minutes.

INSTRUCTOR COMPLIANCE

- Recommend that a certified REPULS Instructor performs this procedure and reviews the Safety Data Sheet prior to any exposure.
- **Contact Crotega Safety Solutions directly with any questions or concerns.**

Questions? Contact Crotega Safety Solutions • www.crotega.com
(612)-547-3207 • contact@crotega.com